

Macro-moths of Bankhead Moss

S.T. Buckland

Macro-moths of Bankhead Moss

S.T. Buckland

I have included images of all the species that I have recorded at Bankhead Moss. Most of these images were taken at Bankhead Moss, but some were taken elsewhere, to illustrate those species that I had recorded but not photographed. My thanks go to Nigel Voaden for identifying the more difficult species.

Steve Buckland steve@st-andrews.ac.uk 5/May/2024

Contents

Angle shades Antler moth	5
Autumnal rustic	7
Barred straw	8
Beautiful golden Y	9
Bordered white	10
Bright-line brown-eye	11
Brimstone moth	12
Brown rustic	13
Brown silver-line	14
Brown-spot pinion	15
Buff footman	16
Buff-tip	17
Burnished brass	18
Canary-shouldered thorn	19
Centre-barred sallow	20
Chestnut	21
Clouded border	22
Clouded-bordered brindle	23
Clouded drab	24
Clouded silver	25
Common carpet	26
Common lutestring	27
Common marbled carpet	28

5	Common pug	29
6	Common quaker	30
7	Common rustic	31
	Common wainscot	32
8	Common wave	33
9	Common white wave	34
10	Coxcomb prominent	35
11	Currant pug	36
12		
13	Dark arches	37
14	Dark marbled carpet	38
15	Dotted border	39
16	Dotted clay	40
17	Double lobed	41
18	Double-striped pug	42
	Dun-bar	43
19		
20	Early thorn	44
21	Engrailed	45
22		
23	Flame carpet	46
24	Flame shoulder	47
25		
26	Garden tiger	48
27	Ghost moth	49
28	Gold swift	50
	Golden-rod pug	51
	Green arches	52
	Green carpet	53
	Grey pug	54

~ ~		
29	Heart and dart	55
30	Hebrew character	56
31		
32	Ingrailed clay	57
33	Iron prominent	58
34	-	
35	July highflyer	59
36		
	Knot grass	60
37		
38	Large emerald	61
39	Large yellow underwing	62
40	Latticed heath	63
41	Lempke's gold spot	64
42	Lesser broad-bordered	
43	yellow underwing	65
	Lesser swallow prominent	66
44	Lesser yellow underwing	67
45	Light emerald	68
	Lunar thorn	69
46	Lunar underwing	70
47		
48		
49		
50		

Contents

Map-winged swift March moth Middle-barred minor Mottled beauty Mottled grey Mottled pug Mouse moth
Nut-tree tussock
Orange sallow
Pale brindled beauty Pale prominent Pale-shouldered brocade Peach blossom Pebble prominent Peppered moth Phoenix Pinion-streaked snout Pink-barred sallow Poplar hawk-moth Purple clay
Red-green carpet Red-necked footman Red sword-grass Riband wave

Rustic

71	Satellite	96
72	Satin beauty	97
73	Scalloped hazel	98
74	Setaceous hebrew characte	r 99
75	Shaded broad-bar	100
76	Shoulder stripe	101
77	Shuttle-shaped dart	102
	Silver-ground carpet	103
78	Silver Y	104
	Single-dotted wave	105
79	Six-striped rustic	106
	Slender brindle	107
80	Small angle shades	108
81	Small dotted buff	109
82	Small fan-footed wave	110
83	Small phoenix	111
84	Small quaker	112
85	Small rivulet	113
86	Small square-spot	114
87	Small wainscot	115
88	Smoky wainscot	116
89	Snout	117
90	Spectacle	118
	Spruce carpet	119
91	Square-spot rustic	120
92	Straw dot	121
93	Suspected	122
94		
05		

,	Tawny-barred angle Twin-spotted quaker	123 124
)	V-pug	125
	Water carpet White ermine Winter moth	126 127 128
	Yellow-horned	129



Angle shades

Common across a range of habitats and through most of the year. Autumn numbers are boosted by immigrants.





Antler moth

A common species with antler-like markings, often seen feeding on flowers such as ragwort during the day.





Autumnal rustic

A common autumn species of open country.





Barred straw

Common in a variety of lowland habitats.





Beautiful golden Y

A fairly common species in a range of habitats.





Bordered white

Common where pines grow.





Bright-line brown-eye

A summer species that occurs in a wide range of habitats.





Brimstone moth

A common woodland species.





Brown rustic

A common species of broadleaved woodland, and occurs in a range of other habitats.



Brown silver-line

A common species of early summer, wherever bracken occurs.





Brown-spot pinion

A declining autumn species, but still quite common in Fife.





Buff footman

This species is a recent arrival in Fife, spreading from the south.





Buff-tip

This moth of open woodland is uncommon in Fife. When at rest, it looks like a broken birch twig.





Burnished brass

A common species, especially anywhere that nettles grow.





Canary-shouldered thorn

A common woodland species in autumn.



Centre-barred sallow

A common autumn species where ashes grow.





Chestnut

Chestnuts overwinter as adults and are usually seen in autumn and early spring.





Clouded border

A common woodland species.





Clouded-bordered brindle

A common species of broadleaved woodland.





Clouded drab

A common woodland species in spring. Its markings are variable.





Clouded silver

A woodland species that is expanding its range in Scotland.





Common carpet

A common summer species in a range of habitats.





Common lutestring

A sparsely distributed species in east Fife. There are plenty of birches to provide food for the caterpillars at Bankhead Moss.





Common marbled carpet

A variable species that is common in summer and autumn in a range of habitats.



Common pug

A common summer pug in a range of habitats.



Common quaker

A very common spring moth.





Common rustic / lesser common rustic

Two common grassland species with variable markings that cannot be separated from images.



Common wainscot

Very similar to the more common smoky wainscot, but with white underwings instead of smoky.



Common wave

A common species of damp woodland, especially where sallows or poplars grow.





Common white wave

A common woodland species, especially where birches, alders or sallows grow.





Coxcomb prominent

A fairly common woodland species.





Currant pug

Occurs where wild hops or garden currants grow.



Dark arches

A common species of grasslands, and wherever grasses grow.





Dark marbled carpet

A common species of woodland and moorland.





Dotted border

A common spring species, occurring mostly in woodland.





Dotted clay

A common summer species in a range of habitats.





Double lobed

This species has expanded its range, but is still very scarce in Fife. This worn individual was at Bankhead Moss in early August 2023.





Double-striped pug

A widely-distributed species, occurring in a variety of habitats. This one has one hind wing that did not expand when it emerged from a pupa.





Dun-bar

A common species, especially in woodland.





Early thorn

This attractive moth is common in woodland in late spring.





Engrailed

A fairly common woodland species, mostly of late spring and early summer.





Flame carpet

A common species across a range of habitats.



Flame shoulder

A common species across a range of habitats.



Garden tiger

Common in open habitats but declining.





Ghost moth

A fairly common grassland species. Female above, male below.





Gold swift

A locally-distributed species, but can be quite abundant where it occurs. It is often associated with bracken.





Golden-rod pug

A fairly common species that occurs in a range of habitats.





Green arches

A large, impressive species of broadleaved woodland.





Green carpet

A common species that occurs in a range of habitats.





Grey pug

A common species that occurs in a range of habitats.





Heart and dart

Common in a very wide range of habitats.



Hebrew character

A very common spring species across a wide range of habitats.





Ingrailed clay

A fairly common and very variable species of woodland and moorland.









Iron prominent

A woodland species of late spring and early summer that is becoming more common.







July highflyer

Common in a range of habitats.



Knot grass

A species of open habitats that is fairly scarce in Fife.





Large emerald

A fairly common and impressive woodland species.





Large yellow underwing

A large, variable and very common grassland species.





Latticed heath

This species is widely-distributed but not especially common in Fife.





Lempke's gold spot

A locally-distributed species of damp habitats. This one was at Bankhead Moss in early July.



Lesser broad-bordered yellow underwing

Common and sometimes abundant across a range of habitats.



Lesser swallow prominent

A fairly common species where birch trees grow.





Lesser yellow underwing

This variable species is common across a range of habitats.





Light emerald

A common species, especially in broadleaved woodland.



68



Lunar thorn

A locally-distributed woodland species.





Lunar underwing

A common autumn species in grassland habitats.





Map-winged swift

A common species in areas of Fife with bracken.





March moth

A common species from mid-February to mid-April, especially in open woodland habitats.





Middle-barred minor

A common summer species of grasslands, often seen on flowerheads in daylight.





Mottled beauty

A common species, especially in woodland.



Mottled grey

An early spring species, occurring in small numbers in mostly open habitats. These were at Bankhead Moss in mid-March 2024.





Mottled pug

A local species of woodland and scrub in Fife.





Mouse moth

A common and widespread species. When threatened, it scuttles like a mouse, hence its name.





Nut-tree tussock

A common species of broadleaved woodland.





Orange sallow

A scarce species of deciduous woodland in Fife. This one was at Bankhead Moss in early September 2023.





Pale brindled beauty

A common species of broadleaved woodland in February and early March.





Pale prominent

A locally-distributed species, occurring where poplars grow.





Pale-shouldered brocade

A common woodland species.





Peach blossom

A scarce species in Fife. Although a big, showy moth (above), it is remarkably difficult to spot when it folds its wings around a twig (below).





Pebble prominent

Occurs where sallows or poplars grow.



Peppered moth

A fairly common species in woodland.



Phoenix

A scarce species in Fife, occurring in gardens and woodland. Its foodplants are black- and redcurrants and goose berries.



Pinion-streaked snout

A scarce species of damp habitats. This one was at Bankhead Moss in late September 2023.



Pink-barred sallow

A common autumn species, especially in broad-leaved woodland and marshes.





Poplar hawkmoth

The most widely-distributed and common hawkmoth in Fife.





Purple clay

A fairly common species of broadleaved woodland and wooded heathland.



Red-green carpet

A common autumn species, and again in spring, when it emerges from hibernation.









Red-necked footman

A scarce woodland species in Fife. These were at Bankhead Moss in late June and early July 2023.





Red sword-grass

A widely-distributed but scarce species of moorland, rough pasture and damp woodland. The one above was at Bankhead Moss in early April 2023, and the one below in mid-March 2024.





Riband wave

Common across a range of habitats.





Rustic

A common species in lowland habitats.





Satellite

A common species in autumn and spring, mostly in broadleaved woodland.





Satin beauty

This species is most common in conifer plantations, but also occurs in ancient broadleaved woodland.





Scalloped hazel

A common woodland species of late spring and early summer.





Setaceous hebrew character

A common lowland species.





Shaded broad-bar

Common in grassy habitats, and often disturbed during the day.



Shoulder stripe

A spring species that occurs where dog roses grow, mostly in woodland and along hedgerows.



Shuttle-shaped dart

A southern species that has spread into Scotland, occurring in a range of habitats.



Silver-ground carpet

A common and often abundant species in a range of habitats. Often seen in daylight.



103



Silver Y

An abundant migrant moth, occurring in a wide range of habitats.





Single-dotted wave

This species is close to the northern limit of its range in Fife. It occurs in woodland and damp habitats.





Six-striped rustic

A common species of wet and damp habitats.



Slender brindle

A fairly common woodland species.





Small angle shades

A fairly common species of woodland, moorland and gardens.





Small dotted buff

A species of wet grassland. Not numerous in the east of Fife.





Small fan-footed wave

A common species, mostly in broadleaved woodland.





Small phoenix

A common species, especially in woodland.





Small quaker

A spring species mostly of broadleaved woodland. It is not numerous in Fife. This one was at Bankhead Moss in late March 2024.





Small rivulet

Very similar to the rivulet. It is common, occurring in a range of habitats.





Small square-spot

A common summer species in a range of habitats.





Small wainscot

A locally-distributed species of damp habitats.





Smoky wainscot

A common grassland species.





Snout

A common species where nettles grow.



Spectacle

This species is not numerous in east Fife. The foodplant of its caterpillars is nettles.





Spruce carpet

A common species where conifers grow.



Square-spot rustic

A common grassland species.





Straw dot

A fairly common grassland species.



Suspected

A scarce species. These rather worn individuals were at Bankhead Moss in mid-August 2023.





Tawny-barred angle

Common, occurring mostly in conifer plantations.





Twin-spotted quaker

A species of broadleaved woodlands that has spread north in recent years, and is now common in Fife.





V-pug

This species is becoming more common and widespread. It occurs in a variety of habitats.





Water carpet

A common spring species, occurring mostly in April and May.





White ermine

Common in a wide range of habitats.



Winter moth

A widespread and common species of late autumn and early winter.





Yellow horned

An early spring species of birch woodland.

